arXiv:1010.0584v1 [quant-ph] 25 Sep 2010

Wigner function evolution in self-Kerr Medium derived by Entangled state representation

Li-yun Hu, Zheng-lu Duan, Xue-xiang Xu, and Zi-sheng Wang
College of Physics & Communication Electronics,
Jiangxi Normal University, Nanchang 330022, China

(Dated:)

By introducing the thermo entangled state representation, we convert the calculation of Wigner function (WF) of density operator to an overlap between "two pure" states in a two-mode enlarged Fock space. Furthermore, we derive a new WF evolution formula of any initial state in self-Kerr Medium with photon loss and find that the photon number distribution for any initial state is independent of the coupling factor with Kerr Medium, where the number state is not affected by the Kerr nonlinearity and evolves into a density operator of binomial distribution.

Keywords: Wigner function, Kerr Medium, entangled state representation

PACS numbers:

I. INTRODUCTION

Nonclassicality of optical fields has been a topic of great interest in quantum optics and quantum information processing [1], which is usually associated with quantum interference and entanglement. The phase space Wigner function (WF) [2, 3] of quantum states of light is a powerful tool for investigating such nonclassical effects. The WF was first introduced by Wigner in 1932 to calculate quantum corrections to a classical distribution function of a quantum-mechanical system. The partial negativity of the WF is indeed a good indication of the highly nonclassical character of the state [4] and monitors a decoherence process of a quantum state, e. g. the excited coherent state in both photon-loss and thermal channels [5, 6], the single-photon subtracted squeezed vacuum state in both amplitude decay and phase damping channels [7], and so on [8–12].

Nonlinear interaction of light in a medium provides a very useful framework to study various nonclassical properties of quantum states of radiation. The Kerr medium is one of the simplest nonlinearity, which shall allow us to investigate the full time-dependent WF dynamics with or without a quantum noise. Recently, a Fokker-Planck equation for the WF evolution in a noisy Kerr medium ($\chi^{(3)}$ nonlinearity) is presented [13]. Then the authors numerically solved this equation assuming coherent state as an initial condition and discussed its dissipation effects. However, for any initial condition, as far as we are concerned, there is no report about the WF evolution. On the other hand, by using the thermo entangled state representation (TESR) we solved various master equations to obtain density operators with an infinite operator-sum representation [14] and then revealed that the WF of density operator can be expressed as an overlap between two pure states (see Eq. (20) below) [15]. This brings much convenience to calculate time evolution of WFs when quantum decoherence happens. Thus the TESR is beneficial to quantum decoherence theory.

In this paper, we shall appeal the TESR $|\eta\rangle$ to treat the WF evolution at any initial condition in self-Kerr Medium with photon loss and present a new formula to calculate time evolution of the WF for quantum decoherence. In addition, based on the derived WF evolution formula, we shall deduce the photon number distribution for any initial state in presence of Kerr interaction, where the photon number distribution is independent of the coupling factor χ that is relative to the Kerr medium. As examples, the WF formula is applied to the cases of initial coherent state, and number state, respectively. Conclusions are involved in the last section.

II. BRIEF REVIEW OF THERMO ENTANGLED STATE REPRESENTATION

We begin with briefly reviewing the thermo entangled state representation (TESR). On the basis of Umezawa-Takahash thermo field dynamics (TFD) [16–18] we constructed the TESR in the doubled Fock space [19–21],

$$\begin{aligned} |\eta\rangle &= D(\eta)\,|\eta=0\rangle \\ &= \exp\left[-\frac{1}{2}|\eta|^2+\eta a^\dagger-\eta^*\tilde{a}^\dagger+a^\dagger\tilde{a}^\dagger\right]\left|0,\tilde{0}\right\rangle, \end{aligned} \tag{1}$$

where $D(\eta)=e^{\eta a^\dagger-\eta^* a}$ is the displacement operator, $\tilde{a}^\dagger$ is a fictitious mode accompanying the real photon creation operator $a^\dagger$, $\left|0,\tilde{0}\right\rangle=|0\rangle\left|\tilde{0}\right\rangle$, and $\left|\tilde{0}\right\rangle$ is annihilated by $\tilde{a}$ with the relations $\left[\tilde{a},\tilde{a}^\dagger\right]=1$ and $\left[a,\tilde{a}^\dagger\right]=0$. The structure of $|\eta\rangle$ is similar to that of the EPR eigenstate shown in Ref. [19]. Operating a and $\tilde{a}$ on $|\eta\rangle$ in Eq.(1) we can obtain the eigen-equations of $|\eta\rangle$,

$$\begin{aligned} &(a-\tilde{a}^\dagger)\,|\eta\rangle=\eta\,|\eta\rangle\,,\ (a^\dagger-\tilde{a})\,|\eta\rangle=\eta^*\,|\eta\rangle\,, \\ &\langle\eta|\,(a^\dagger-\tilde{a})=\eta^*\,\langle\eta|\,,\ \langle\eta|\,(a-\tilde{a}^\dagger)=\eta\,\langle\eta|\,. \end{aligned} \tag{2}$$

Note that $\left[(a-\tilde{a}^\dagger),(a^\dagger-\tilde{a})\right]=0$, thus $|\eta\rangle$ is the common eigenvector of $(a-\tilde{a}^\dagger)$ and $(\tilde{a}-a^\dagger)$. Using the normally ordered form of vacuum projector $\left|0,\tilde{0}\right\rangle\left\langle 0,\tilde{0}\right|=\colon\exp\left(-a^\dagger a-\tilde{a}^\dagger\tilde{a}\right)\colon$ and the technique of integration within an ordered product (IWOP) of operators [22–24], we can easily prove that $|\eta\rangle$ is complete and orthonormal,

$$\int \frac{\mathrm{d}^2\eta}{\pi} |\eta\rangle \langle\eta| = 1, \;\; \langle\eta' |\eta\rangle = \pi\delta\left(\eta' - \eta\right)\delta\left(\eta'^{*} - \eta^{*}\right). \tag{3}$$

It is easily seen that $|\eta = 0\rangle$ has the properties

$$|\eta = 0\rangle = e^{a^\dagger \tilde{a}^\dagger} \left|0, \tilde{0}\right\rangle = \sum_{n=0}^{\infty} |n, \tilde{n}\rangle, \tag{4}$$

(where $n = \tilde{n}$, and $\tilde{n}$ denotes the number in the fictitious Hilbert space) and

$$\begin{aligned} a \;\; |\eta = 0\rangle &= \tilde{a}^\dagger |\eta = 0\rangle, \\ a^\dagger |\eta = 0\rangle &= \tilde{a} |\eta = 0\rangle, \\ \left(a^\dagger a\right)^n |\eta = 0\rangle &= \left(\tilde{a}^\dagger \tilde{a}\right)^n |\eta = 0\rangle. \end{aligned} \tag{5}$$

Note that density operators $\rho(a^\dagger, a)$ are defined in the real space which are commutative with operators $(\tilde{a}^\dagger, \tilde{a})$ in the tilde space.

In a similar way, we can introduce the state vector $|\xi\rangle$ conjugated to $|\eta\rangle$, defined as

$$\begin{aligned} |\xi\rangle &= D\left(\xi\right) e^{-a^\dagger \tilde{a}^\dagger} \left|0, \tilde{0}\right\rangle \\ &= \exp\left(-\frac{1}{2}|\xi|^2 + \xi a^\dagger + \xi^{*}\tilde{a}^\dagger - a^\dagger \tilde{a}^\dagger\right) \left|0, \tilde{0}\right\rangle \\ &= (-1)^{a^\dagger a} |\eta = -\xi\rangle, \end{aligned} \tag{6}$$

which also possesses orthonormal and complete properties

$$\int \frac{\mathrm{d}^2\xi}{\pi} |\xi\rangle \langle\xi| = 1, \;\; \langle\xi' |\xi\rangle = \pi\delta\left(\xi' - \xi\right)\delta\left(\xi'^{*} - \xi^{*}\right). \tag{7}$$

III. MASTER EQUATION FOR A SELF-KERR INTERACTION

In the Markov approximation and interaction picture the master equation for a dissipative cavity with Kerr medium has the form [25, 26]

$$\frac{d\rho}{dt} = -i\chi \left[\left(a^\dagger a\right)^2, \rho\right] + \gamma\left(2a\rho a^\dagger - a^\dagger a\rho - \rho a^\dagger a\right), \tag{8}$$

where γ is decaying parameter of the dissipative cavity, χ is coupling factor depending on the Kerr medium. Milburn and Holmes [27] solved this equation by changing it to a partial differential equation for the Q-function and for an initial coherent state. Here we will solve the master equation by virtue of the entangled state representation and present the infinite sum representation of density operator.

Operating the both sides of Eq.(8) on the state $|\eta = 0\rangle$, letting $|\rho\rangle = \rho|\eta = 0\rangle$ (Here one should understand the single-mode density operator ρ in the left of Eq.(8) as the direct product $\rho \otimes \tilde{I}$ when ρ acts onto the two-mode state $|\eta = 0\rangle = e^{a^\dagger \tilde{a}^\dagger} \left|0, \tilde{0}\right\rangle$, where $\tilde{I}$ is the identity operator in the auxiliary mode.), and using Eq.(5) we can convert the master equation in Eq. (8) into the following form,

$$\begin{aligned} \frac{d}{dt}|\rho\rangle \;=\; & \left\{-i\chi\left[\left(a^\dagger a\right)^2 - \left(\tilde{a}^\dagger \tilde{a}\right)^2\right]\right. \\ & \left. + \gamma\left(2a\tilde{a} - a^\dagger a - \tilde{a}^\dagger \tilde{a}\right)\right\} |\rho\rangle, \end{aligned} \tag{9}$$

i.e., an evolution equation of state vector $|\rho\rangle$. Its solution is then of the form

$$|\rho\rangle = e^{-i\chi t\left[\left(a^\dagger a\right)^2 - \left(\tilde{a}^\dagger \tilde{a}\right)^2\right] + \gamma t\left(2a\tilde{a} - a^\dagger a - \tilde{a}^\dagger \tilde{a}\right)} |\rho_0\rangle, \tag{10}$$

where $|\rho_0\rangle = \rho_0 |\eta = 0\rangle$, ρ_0 is an initial density operator. The advantage of using thermo field notation over more traditional algebraic manipulation with superoperators is that in many situations (and, particularly, ones of our interest) it enables to simplify, make more illustrative and less cumbersome finding the solution (10) and estimation of time-dependent matrix elements. In particular, it allows to represent in a simple form a factorization of the superoperator $\exp\{\cdots\}$ into multipliers with easily estimated actions on the number states [28].

By introducing the following operators,

$$K_0 = a^\dagger a - \tilde{a}^\dagger \tilde{a}, \;\; K_z = \frac{a^\dagger a + \tilde{a}^\dagger \tilde{a} + 1}{2}, \;\; K_- = a\tilde{a}, \tag{11}$$

which satisfy $[K_0, K_z] = [K_0, K_-] = 0$, we can rewrite Eq.(10) as

$$\begin{aligned} |\rho\rangle &= e^{\{-i\chi t[K_0(2K_z - 1)] + \gamma t(2K_- - 2K_z + 1)\}} |\rho_0\rangle \\ &= \exp\left[i\chi t K_0 + \gamma t\right] \\ &\times \exp\left\{-2t\left(\gamma + i\chi K_0\right)\left[K_z + \frac{-\gamma}{\gamma + i\chi K_0} K_-\right]\right\} |\rho_0\rangle. \end{aligned} \tag{12}$$

With the aid of the operator identity [30]

$$\begin{aligned} e^{\lambda(A + \sigma B)} \;&=\; e^{\lambda A} \exp\left[\sigma B\left(1 - e^{-\lambda\tau}\right)/\tau\right] \\ &=\; \exp\left[\sigma B\left(e^{\lambda\tau} - 1\right)/\tau\right] e^{\lambda A}, \end{aligned} \tag{13}$$

which is valid for $[A, B] = \tau B$, and noticing $[K_z, K_-] = -K_-$, we can reform Eq.(12) as

$$|\rho\rangle = \exp\left[i\chi t K_0 + \gamma t\right] \exp\left[\Gamma_z K_z\right] \exp\left[\Gamma_- K_-\right] |\rho_0\rangle, \tag{14}$$

where

$$\Gamma_z = -2t\left(\gamma + i\chi K_0\right), \;\; \Gamma_- = \frac{\gamma\left(1 - e^{-2t(\gamma + i\chi K_0)}\right)}{\gamma + i\chi K_0}. \tag{15}$$

From Eq.(14) we can obtain the infinite operator-sum form of $\rho\left(t\right)$, (see Appendix A)

$$\rho\left(t\right) = \sum_{m,n,l=0}^{\infty} M_{m,n,l} \rho_0 \mathcal{M}^\dagger_{m,n,l}, \tag{16}$$

where the two operators $M_{m,n,l}$ and $\mathcal{M}^{\dagger}_{m,n,l}$ are respectively defined as

$$M_{m,n,l} \equiv \sqrt{\frac{\Lambda^{l}_{m,n}}{l!}} e^{-i\chi t m^{2}-\gamma t m} \left|m\right\rangle \left\langle m\right| a^{l},$$

$$\mathcal{M}^{\dagger}_{m,n,l} \equiv \left\{ \sqrt{\frac{\Lambda^{l}_{n,m}}{l!}} e^{-i\chi t n^{2}-\gamma t n} \left|n\right\rangle \left\langle n\right| a^{l} \right\}^{\dagger}. \qquad (17)$$

Although $M_{m,n,l}$ is not hermite conjugate to $\mathcal{M}^{\dagger}_{m,n,l}$, the normalization still holds, ($\sum_{m,n,l=0}^{\infty}\mathcal{M}^{\dagger}_{m,n,l}M_{m,n,l} = 1$, see Appendix B [29]) i.e., they are trace-preserving in a general sense, so $M_{m,n,l}$ and $\mathcal{M}^{\dagger}_{m,n,l}$ may be named the generalized Kraus operators.

IV. EVOLUTION OF WIGNER FUNCTION FOR SELF-KERR CHANNEL

In this section, we consider Wigner function's time evolution in the self-Kerr medium channel. For this purpose, we shall derive a new expression of Wigner function in the TESR. According to the definition of Wigner function of density operator ρ,

$$W\left(\alpha,\alpha^{*}\right) = \mathrm{Tr}\left[\Delta\left(\alpha,\alpha^{*}\right)\rho\right], \qquad (18)$$

where $\Delta\left(\alpha,\alpha^{*}\right)$ is the single-mode Wigner operator [2, 30], whose explicit normally ordered form is [31]

$$\Delta\left(\alpha,\alpha^{*}\right) = \frac{1}{\pi} :e^{-2\left(a^{\dagger}-\alpha^{*}\right)\left(a-\alpha\right)}: = \frac{1}{\pi} D\left(2\alpha\right)\left(-1\right)^{a^{\dagger}a}. \qquad (19)$$

By using $\left\langle\tilde{n}\right|\left.\tilde{m}\right\rangle = \delta_{n,m}$ ($n=\tilde{n}, m=\tilde{m}$) and noticing (6) as well as $\left|\rho\right\rangle = \rho\left|\eta=0\right\rangle$, we can reform Eq.(18) as

$$\begin{aligned} W\left(\alpha,\alpha^{*}\right) &= \sum_{m,n}^{\infty}\left\langle n,\tilde{n}\right|\Delta\left(\alpha,\alpha^{*}\right)\rho\left|m,\tilde{m}\right\rangle \\ &= \frac{1}{\pi}\left\langle\eta=0\right|D\left(2\alpha\right)\left(-1\right)^{a^{\dagger}a}\left|\rho\right\rangle \\ &= \frac{1}{\pi}\left\langle\eta=-2\alpha\right|\left(-1\right)^{a^{\dagger}a}\left|\rho\right\rangle \\ &= \frac{1}{\pi}\left\langle\xi=2\alpha\right|\left.\rho\right\rangle, \end{aligned} \qquad (20)$$

where Eq. (20) is the Wigner function formula in thermo entangled state representation, with which the Wigner function of density operator is simplified as an overlap between two "pure states" in enlarged Fock space, rather than using ensemble average in the system-mode space. This will brings much convenience to calculate the time evolution of Wigner functions when quantum decoherence happens.

Projecting (14) on the entangled state representation $\frac{1}{\pi}\left\langle\xi_{=2\alpha}\right|$, and inserting the completeness relation (7), we find

$$W\left(\alpha,\alpha^{*},t\right) = 4\int\frac{\mathrm{d}^{2}\beta}{\pi}G\left(\alpha,\beta,t\right)W\left(\beta,\beta^{*},0\right), \qquad (21)$$

where $W\left(\alpha,\alpha^{*},t\right)$ and $W\left(\beta,\beta^{*},0\right)$ are the Wigner functions at the evolving time t and initial time, respectively, and

$$\begin{aligned} G\left(\alpha,\beta,t\right) &= \left\langle\xi_{=2\alpha}\right|\exp\left[i\chi tK_{0}+\gamma t\right] \\ &\quad\times\exp\left[\Gamma_{z}K_{z}\right]\exp\left[\Gamma_{-}K_{-}\right]\left|\xi'_{=2\beta}\right\rangle. \end{aligned} \qquad (22)$$

It is convenient to the matrix element in (22) according to the two-mode Fock space. Thus the $\left\langle\xi_{=2\alpha}\right|$ is expanded as

$$\left\langle\xi\right| = \left\langle0,\tilde{0}\right|\sum_{m,n=0}^{\infty}\frac{a^{m}\tilde{a}^{n}}{m!n!}H_{m,n}\left(\xi^{*},\xi\right)e^{-\left|\xi\right|^{2}/2}. \qquad (23)$$

By using the two-mode Fock state $\left|m,\tilde{n}\right\rangle = a^{\dagger m}\tilde{a}^{\dagger n}/\sqrt{m!n!}\left|0,\tilde{0}\right\rangle$, we get

$$\left\langle\xi\right|\left.m,\tilde{n}\right\rangle = H_{m,n}\left(\xi^{*},\xi\right)e^{-\left|\xi\right|^{2}/2}/\sqrt{m!n!}, \qquad (24)$$

where $H_{m,n}\left(\xi^{*},\xi\right)$ is the two-variable Hermite polynomials [32, 33]. Inserting the complete relation $\sum_{m,n=0}^{\infty}\left|m,\tilde{n}\right\rangle\left\langle m,\tilde{n}\right| = 1$, after a long but straight calculation, then the Wigner function's evolution is given by (see Appendix C)

$$W\left(\alpha,\alpha^{*},t\right) = \sum_{m,n=0}^{\infty}C_{m,n}\left(\alpha,\alpha^{*},t\right)E_{m,n}, \qquad (25)$$

where $\Lambda_{m,n} \equiv \frac{\gamma\left(1-e^{-2t\left(\gamma+i\chi\left(m-n\right)\right)}\right)}{\gamma+i\chi\left(m-n\right)}$,

$$\begin{aligned} &C_{m,n}\left(\alpha,\alpha^{*},t\right) \\ &\equiv \frac{e^{-i\chi t\left(m^{2}-n^{2}\right)-\gamma t\left(m+n\right)}e^{-2\left|\alpha\right|^{2}}}{m!n!\left(\Lambda_{m,n}+1\right)^{\left(m+n+2\right)/2}}H_{m,n}\left(2\alpha^{*},2\alpha\right), \end{aligned} \qquad (26)$$

and

$$\begin{aligned} E_{m,n} &= 4\int\frac{\mathrm{d}^{2}\beta}{\pi}W\left(\beta,\beta^{*},0\right)e^{\frac{2\left(\Lambda_{m,n}-1\right)}{\Lambda_{m,n}+1}\left|\beta\right|^{2}} \\ &\quad\times H_{m,n}\left(\frac{2\beta}{\sqrt{\Lambda_{m,n}+1}},\frac{2\beta^{*}}{\sqrt{\Lambda_{m,n}+1}}\right). \end{aligned} \qquad (27)$$

It is obvious that, when $\chi=0$, the case of photon loss, $\Lambda_{m,n}\rightarrow\left(1-e^{-2\gamma t}\right)=T$ and Eq.(25) just does reduce to (see Appendix E)

$$W\left(\alpha,\alpha^{*},t\right) = \frac{2}{T}\int\frac{\mathrm{d}^{2}\beta}{\pi}e^{-\frac{2}{T}\left|\alpha-\beta e^{-\gamma t}\right|^{2}}W\left(\beta,\beta^{*},0\right), \qquad (28)$$

which is just the evolving formula of Wigner function for amplitude-damping channel. While for $\gamma=0$, without photon-loss, Eq.(25) reduces to

$$\begin{aligned} &W\left(\alpha,\alpha^{*},t\right) \\ &= \sum_{m,n=0}^{\infty}\frac{\exp\left[-i\chi t\left(m^{2}-n^{2}\right)\right]}{m!n!e^{2\left|\alpha\right|^{2}}}H_{m,n}\left(2\alpha^{*},2\alpha\right) \\ &\quad\times 4\int\frac{\mathrm{d}^{2}\beta}{\pi}e^{-2\left|\beta\right|^{2}}H_{m,n}\left(2\beta,2\beta^{*}\right)W\left(\beta,\beta^{*},0\right). \end{aligned} \qquad (29)$$

V. PHOTON NUMBER DISTRIBUTION IN PRESENCE OF KERR INTERACTION

Now we consider photon number (PN) distribution in presence of Kerr medium. According to the TFD, we can reform the PN $p(n)=\mathtt{tr}\left[\rho\left|n\right\rangle\left\langle n\right|\right]$ as

$$\begin{aligned}p(n) &= \left\langle n\right|\rho\left|n\right\rangle=\sum_{m=0}^{\infty}\left\langle n,\tilde{n}\right|\rho\left|m,\tilde{m}\right\rangle \\ &= \left\langle n,\tilde{n}\right|\rho\left|\eta=0\right\rangle=\left\langle n,\tilde{n}\right|\left.\rho\right\rangle, \qquad (30)\end{aligned}$$

which is converted to the matrix element $\left\langle n,\tilde{n}\right|\left.\rho\right\rangle$ in the context of thermo dynamics. Then using the completeness of $\left\langle\xi\right|$ and Eq.(30) as well as Eq.(20), we have

$$\begin{aligned}p(n) &= \int\frac{\mathtt{d}^2\xi}{\pi}\left\langle n,\tilde{n}\right|\left.\xi\right\rangle\left\langle\xi\right|\left.\rho\right\rangle \\ &= \int\mathtt{d}^2\xi\left\langle n,\tilde{n}\right|\left.\xi\right\rangle W\left(\alpha=\xi/2,\alpha^*=\xi^*/2\right) \\ &= 4\pi\int\mathtt{d}^2\alpha W_{\left|n\right\rangle\left\langle n\right|}\left(\alpha,\alpha^*\right)W\left(\alpha,\alpha^*\right), \qquad (31)\end{aligned}$$

where $W_{\left|n\right\rangle\left\langle n\right|}\left(\alpha,\alpha^*\right)=\frac{(-1)^n}{\pi}e^{-2|\alpha|^2}L_n(4\left|\alpha\right|^2)$ is the Wigner function of number state $\left|n\right\rangle\left\langle n\right|$ as shown in [34, 35]. Thus one can calculate the PN by combining Eqs.(25) and (31).

Next we evaluate the PN for the above decoherence model in Eq.(8). Substituting Eq.(25) into Eq.(31), we have

$$\begin{aligned}p(s) &= 4\pi\sum_{m,n=0}^{\infty}\int\mathtt{d}^2\alpha W_{\left|s\right\rangle\left\langle s\right|}\left(\alpha,\alpha^*\right)C_{m,n}\left(\alpha,\alpha^*,t\right)E_{m,n} \\ &= \sum_{m,n=0}^{\infty}\frac{4\pi e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}}{m!n!\left(\Lambda_{m,n}+1\right)^{(m+n+2)/2}}E_{m,n}\cdot F_{m,n}, \qquad (32)\end{aligned}$$

where

$$\begin{aligned}F_{m,n} &\equiv \int\mathtt{d}^2\alpha e^{-2|\alpha|^2}W_{\left|s\right\rangle\left\langle s\right|}\left(\alpha,\alpha^*\right)H_{m,n}\left(2\alpha^*,2\alpha\right) \\ &= \frac{s!}{4}\delta_{m,s}\delta_{n,s}. \qquad (33)\end{aligned}$$

Then substituting Eqs.(33) and (27) into (32) yields

$$\begin{aligned}p(s) &= \frac{4(-1)^s e^{2\gamma t}}{\left(2e^{2\gamma t}-1\right)^{s+1}}\int\mathtt{d}^2\beta\exp\left\{-\frac{2\left|\beta\right|^2}{2e^{2\gamma t}-1}\right\} \\ &\quad\times L_s\left(\frac{4e^{2\gamma t}\left|\beta\right|^2}{2e^{2\gamma t}-1}\right)W\left(\beta,\beta^*,0\right), \qquad (34)\end{aligned}$$

which corresponds to the photon number of density operator in the amplitude-damping quantum channel [15]. From Eq.(34), it is easily to see that, for any initial state, the photon number distribution $p(s)$ is independent of the coupling factor χ that is relative to the Kerr medium, as respected in [36].

VI. EVOLUTION OF QUANTUM STATES

The phase space Wigner distribution function description of quantum states of light is a powerful tool to investigate nonclassical effects, such as quantum interference and entanglement. In this section, as the applications of WF evolution formula, we take two special initial states as examples.

(1) When the initial state is the coherent state $\left|z\right\rangle$, whose WF is given $W\left(\beta,\beta^*\right)=\frac{1}{\pi}e^{-2|\beta-z|^2}$, thus substituting into Eq.(27) yields (see Appendix G)

$$E_{m,n}=\frac{1}{\pi}\left(\Lambda_{m,n}+1\right)^{\frac{m+n+2}{2}}e^{\left(\Lambda_{m,n}-1\right)|z|^2}z^m z^{*n}, \qquad (35)$$

so

$$\begin{aligned}&W\left(\alpha,\alpha^*,t\right) \\ &= \frac{e^{-2|\alpha|^2}}{\pi}\sum_{m,n=0}^{\infty}\frac{z^m z^{*n}}{m!n!}e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)} \\ &\quad\times e^{\left(\Lambda_{m,n}-1\right)|z|^2}H_{m,n}\left(2\alpha^*,2\alpha\right), \qquad (36)\end{aligned}$$

which is a new expression of the evolution of WF for any initial state. In particular, when $\gamma=0$, without the dissipation, Eq.(36) reduces to

$$\begin{aligned}&W\left(\alpha,\alpha^*,t\right) \\ &= \frac{e^{-|z|^2}}{\pi e^{2|\alpha|^2}}\sum_{m,n=0}^{\infty}\frac{z^m z^{*n}}{m!n!}e^{-i\chi t(m^2-n^2)}H_{m,n}\left(2\alpha^*,2\alpha\right), \qquad (37)\end{aligned}$$

which is identical to Eq.(7) in Ref.[13], where Eq.(7) is used to make the numerical calculation since it is much more rapid than the other expression Eq.(6). In addition, from Eq.(36) one can see that the WF can be obtained very quickly when the dissipation cannot be negligible. Further when $\chi t=2\pi$, Eq.(37) just returns to the WF of the initial coherent state.

Fig.1 presents the plots of the WF for different parameters χt and $\alpha=2$. From Fig.1, one can see that the WF turns into an ellipse and squeezing appears in an appropriate direction. Then the ellipse changes into a banana shape and a tailor of the interference fringes appears where the distribution takes the negative values.

(2) Another example is number state, where the WF of number state $\left|s\right\rangle$ is given by

$$\begin{aligned}W_s\left(\beta,\beta^*,0\right) &= \frac{1/s!}{\pi}e^{-2|\beta|^2}H_{s,s}\left(2\beta,2\beta^*\right) \\ &= \frac{(-1)^s}{\pi}e^{-2|\beta|^2}L_s(4\left|\beta\right|^2), \qquad (38)\end{aligned}$$

substituting it into (27) and using the generating function of two-variable Hermite polynomials [see below Eq.(F3)], we have

$$E_{m,n}=\frac{s!}{\pi}\frac{\Lambda_{m,n}^{s-m}\left(\Lambda_{m,n}+1\right)^{m+1}}{(s-m)!}\delta_{m,n}, \qquad (39)$$



FIG. 1: (Color online) The WF for different parameters χt and $\alpha = 2$, (a) $\chi t = 0$, (b) $\chi t = 0.04$, (c) $\chi t = 0.06$, (d) $\chi t = 0.08$, (e) $\chi t = 0.1$, (f) $\chi t = 0.2$.

thus the evolution of WF for $|s\rangle$ is

$$W_s(\alpha,\alpha^*,t) = \sum_{m=0}^{s}\binom{s}{m} e^{-2m\gamma t}\left(1-e^{-2\gamma t}\right)^{s-m} W_m(\alpha,\alpha^*,0) \tag{40}$$

which is the WF of number state in the photon-loss channel and indicates that the number state is not affected by the Kerr nonlinearity. In particular, when $\gamma = 0$, or $t = 0$, Eq.(40) just reduces to the WF of number state. From Eq.(40), on the other hand, it is found that the number state evolves into a density operator of binomial distribution (a mixed state) if $e^{-2\gamma t}$ is a binomial parameter.

VII. CONCLUSIONS

In summary, by converting Wigner function for quantum state into an overlap between two "pure states" in a two-mode enlarged Fock space, we investigate the WF evolution of any initial condition in self-Kerr Medium with photon loss and present a new formula for calculating time evolution of the WF for quantum decoherence. Based on the derived WF evolution formula, in addition, we discuss the photon number distribution for any initial state in presence of Kerr interaction. It is found that the photon number distribution is independent of the coupling factor χ in correlation with the Kerr medium, as expected by people. As applications, furthermore, the two cases of initial coherent state and number state are considered. It is shown that the coherent state can be squeezed due to the presence of Kerr medium, while the number state is not affected by the Kerr nonlinearity and evolves into a density operator of binomial distribution (a mixed state) with $e^{-2\gamma t}$ being a binomial parameter.

ACKNOWLEDGEMENTS Work supported by a grant from the Key Programs Foundation of Ministry of Education of China (No. 210115) and the Research Foundation of the Education Department of Jiangxi Province of China (No. GJJ10097).

Appendix A: Derivation of Eq.(16)

In order to obtain the infinite operator-sum form of $\rho(t)$ from Eq.(14), using the completeness relation of Fock state in the enlarged space $\sum_{m,n=0}^{\infty}|m,\tilde{n}\rangle\langle m,\tilde{n}| = 1$ and noticing $a^{\dagger l}|n\rangle = \sqrt{\frac{(l+n)!}{n!}}|n+l\rangle$, we have

$$\begin{aligned}
|\rho\rangle &= e^{i\chi tK_0+\gamma t}e^{\Gamma_z K_z}e^{\Gamma_- K_-}|\rho_0\rangle \\
&= e^{i\chi tK_0+\gamma t}e^{\Gamma_z K_z}\sum_{m,n=0}^{\infty}|m,\tilde{n}\rangle\langle m,\tilde{n}|e^{\Gamma_- K_-}|\rho_0\rangle \\
&= \sum_{m,n=0}^{\infty} e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)} \\
&\quad\times |m,\tilde{n}\rangle\langle m,\tilde{n}|e^{\Lambda_{m,n}a\tilde{a}}|\rho_0\rangle,
\end{aligned} \tag{A1}$$

where

$$\Lambda_{m,n} = \frac{\gamma(1-e^{-2t(\gamma+i\chi(m-n))})}{\gamma+i\chi(m-n)}. \tag{A2}$$

Furthermore, using the relations

$$\langle n|\,\eta=0\rangle = |\tilde{n}\rangle,\quad |m,\tilde{n}\rangle = |m\rangle\langle n|\,\eta=0\rangle, \tag{A3}$$

we find

$$\begin{aligned}
\langle m,\tilde{n}|a^l\rho_0 a^{\dagger l}|\eta=0\rangle &= \langle m|\langle\tilde{n}|a^l\rho_0 a^{\dagger l}|\eta=0\rangle \\
&= \langle m|a^l\rho_0 a^{\dagger l}(\langle\tilde{n}|\,\eta=0\rangle) \\
&= \langle m|a^l\rho_0 a^{\dagger l}|n\rangle.
\end{aligned} \tag{A4}$$

Thus Eq.(A1) becomes

$$\begin{aligned}
|\rho\rangle &= \sum_{m,n,l=0}^{\infty}\frac{\Lambda_{m,n}^l}{l!}e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)} \\
&\quad\times|m,\tilde{n}\rangle\langle m|a^l\rho_0 a^{\dagger l}|n\rangle \\
&= \sum_{m,n,l=0}^{\infty}\frac{\sqrt{(n+l)!\,(m+l)!}}{\sqrt{n!m!}\,l!\,\Lambda_{m,n}^{-l}} \\
&\quad\times e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}|m,\tilde{n}\rangle\rho_{0,m+l,n+l},
\end{aligned} \tag{A5}$$

where $\rho_{0,m+l,n+l} \equiv \langle m+l|\,\rho_0\,|n+l\rangle$. Using Eq.(A3) again, we see

$$|\rho\rangle = \sum_{m,n,l=0}^{\infty} \frac{\sqrt{(n+l)!\,(m+l)!}}{\sqrt{n!m!}\,l!\Lambda_{m,n}^{-l}} \times e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}\rho_{0,m+l,n+l}\,|m\rangle\,\langle n|\;\eta=0\rangle\,. \tag{A6}$$

After depriving $|\eta=0\rangle$ from the both sides of Eq.(A6), the solution of master equation (8) appears as an infinite operator-sum form

$$\begin{aligned}\rho(t) &= \sum_{m,n,l=0}^{\infty} \sqrt{\frac{(n+l)!\,(m+l)!}{n!m!}}\frac{\Lambda_{m,n}^{l}}{l!}e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}\,|m\rangle\,\langle m+l|\,\rho_0\,|n+l\rangle\,\langle n| \\ &= \sum_{m,n,l=0}^{\infty} \frac{\Lambda_{m,n}^{l}}{l!}e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}\,|m\rangle\,\langle m|\,a^l\rho_0 a^{\dagger l}\,|n\rangle\,\langle n|\,.\end{aligned} \tag{A7}$$

Note that the factor $(m-n)$ appears in the denominator of $\Lambda_{m,n}$ (see Eq.(A2)), (this is originated from the nonlinear term of $\left(a^{\dagger}a\right)^2$) so that it is impossible to move all $n-$dependent terms to the right of $a^l\rho_0 a^{\dagger l}$. Fortunately, we can formally express Eq.(A7) as Eq.(16).

Appendix B: Proof of normalization for the generalized Kraus operators

Using the operator identity $e^{\lambda a^{\dagger}a} =\colon \exp\left[\left(e^{\lambda}-1\right)a^{\dagger}a\right]\colon$ and the IWOP technique, we can prove that

$$\begin{aligned}&\sum_{m,n,l=0}^{\infty}\mathcal{M}_{m,n,l}^{\dagger}M_{m,n,l} \\ &= \sum_{n,l=0}^{\infty}\frac{(n+l)!}{n!}\frac{\left(1-e^{-2t\gamma}\right)^l}{l!}e^{-2n\gamma t}\,|n+l\rangle\,\langle n+l| \\ &= \sum_{n,l=0}^{\infty}\frac{\left(1-e^{-2t\gamma}\right)^l}{l!}a^{\dagger l}e^{-2\gamma t a^{\dagger}a}\,|n\rangle\,\langle n|\,a^l \\ &= \sum_{l=0}^{\infty}\frac{\left(1-e^{-2t\gamma}\right)^l}{l!}\colon\exp\left[\left(e^{-2\gamma t}-1\right)a^{\dagger}a\right]\left(a^{\dagger}a\right)^l\colon = 1,\end{aligned} \tag{B1}$$

from which one can see that the normalization still holds, i.e., they are trace-preserving in a general sense, so $M_{m,n,l}$ and $\mathcal{M}_{m,n,l}^{\dagger}$ may be named the generalized Kraus operators.

Appendix C: Derivation of Eq.(25)

Using Eq.(24) and (A1) as well as (A2), Eq.(22) can be rewritten as

$$\begin{aligned}G(\alpha,\beta,t) &= \sum_{m,n=0}^{\infty} e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}\left\langle\xi_{=2\alpha}\,\middle|m,\tilde{n}\right\rangle\left\langle m,\tilde{n}\right|e^{\Lambda_{m,n}a\tilde{a}}\left|\xi'_{=2\beta}\right\rangle \\ &= \sum_{m,n=0}^{\infty}\frac{e^{-2|\alpha|^2}}{\sqrt{m!n!}}e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}H_{m,n}\left(2\alpha^*,2\alpha\right)\sum_{l=0}^{\infty}\frac{\Lambda_{m,n}^{l}}{l!}\left\langle m,\tilde{n}\right|a^l\tilde{a}^l\left|\xi'_{=2\beta}\right\rangle \\ &= e^{-2|\beta|^2-2|\alpha|^2}\sum_{m,n=0}^{\infty}\frac{e^{-i\chi t\left(m^2-n^2\right)-\gamma t(m+n)}}{m!n!}H_{m,n}\left(2\alpha^*,2\alpha\right)\sum_{l=0}^{\infty}\frac{\Lambda_{m,n}^{l}}{l!}H_{m+l,n+l}\left(2\beta,2\beta^*\right).\end{aligned} \tag{C1}$$

Further using a new sum formula (see appendix D)

$$\sum_{l=0}^{\infty}\frac{z^l}{l!}H_{m+l,n+l}\left(x,y\right) = \frac{e^{\frac{zxy}{z+1}}}{(z+1)^{(m+n+2)/2}}H_{m,n}\left(\frac{x}{\sqrt{z+1}},\frac{y}{\sqrt{z+1}}\right), \tag{C2}$$

thus Eq.(C1) can be recast into the following form

$$G\left(\alpha,\beta,t\right) = \sum_{m,n=0}^{\infty}C_{m,n}\left(\alpha,\alpha^*,t\right)e^{2\frac{\Lambda_{m,n}-1}{\Lambda_{m,n}+1}|\beta|^2} \times H_{m,n}\left(\frac{2\beta}{\sqrt{\Lambda_{m,n}+1}},\frac{2\beta^*}{\sqrt{\Lambda_{m,n}+1}}\right), \tag{C3}$$

where $C_{m,n}\left(\alpha,\alpha^*,t\right)$ is defined in (26). Substituting Eq.(C3) into (21) yields (25) and (27).

Appendix D: Derivation of Eq.(C2)

Using the integratal expression of two-mode Hermite polynomials,

$$H_{m,n}\left(\xi,\eta\right) = (-1)^n e^{\xi\eta}\int\frac{d^2z}{\pi}z^n z^{*m}e^{-|z|^2+\xi z-\eta z^*}, \tag{D1}$$

we have

$$\begin{aligned}\sum_{l=0}^{\infty}\frac{\alpha^l}{l!}H_{m+l,n+l}\left(x,y\right) &= \sum_{l=0}^{\infty}\frac{\alpha^l}{l!}(-1)^{n+l}e^{xy}\int\frac{d^2z}{\pi}z^{n+l}z^{*m+l}\exp\left[-\left|z\right|^2+xz-yz^*\right]\\ &= e^{xy}(-1)^n\int\frac{d^2z}{\pi}\sum_{l=0}^{\infty}\frac{\left(-\alpha\left|z\right|^2\right)^l}{l!}z^n z^{*m}\exp\left[-\left|z\right|^2+xz-yz^*\right]\\ &= e^{xy}(-1)^n\int\frac{d^2z}{\pi}z^n z^{*m}\exp\left[-\left(\alpha+1\right)\left|z\right|^2+xz-yz^*\right]\\ &= \frac{e^{\frac{\alpha xy}{\alpha+1}}}{(\alpha+1)^{(m+n+2)/2}}(-1)^n e^{xy/(\alpha+1)}\int\frac{d^2z}{\pi}z^n z^{*m}\exp\left[-\left|z\right|^2+\frac{xz-yz^*}{\sqrt{\alpha+1}}\right]\\ &= \frac{e^{\frac{\alpha xy}{\alpha+1}}}{(\alpha+1)^{(m+n+2)/2}}H_{m,n}\left(\frac{x}{\sqrt{\alpha+1}},\frac{y}{\sqrt{\alpha+1}}\right),\end{aligned} \tag{D2}$$

thus we have completed the proof of (C4).

Appendix E: Derivation of Eq.(28)

When $\chi=0$, $\Lambda_{m,n}\rightarrow(1-e^{-2\gamma t})=T$, and

$$C_{m,n}\left(\alpha,\alpha^*,t\right) \equiv \frac{\exp\left[-\gamma t\left(m+n\right)\right]}{m!n!\left(T+1\right)^{(m+n+2)/2}}H_{m,n}\left(2\alpha^*,2\alpha\right)e^{-2|\alpha|^2}, \tag{E1}$$

then we have

$$\sum_{m,n=0}^{\infty}C_{m,n}\left(\alpha,\alpha^*,t\right)H_{m,n}\left(2\beta,2\beta^*\right) = \frac{e^{-2|\alpha|^2}}{T+1}\sum_{m,n=0}^{\infty}\frac{\left(\frac{e^{-\gamma t}}{\sqrt{T+1}}\right)^{m+n}}{m!n!} \times H_{m,n}\left(2\alpha^*,2\alpha\right)H_{m,n}\left(2\beta,2\beta^*\right). \tag{E2}$$

Using the following formula

$$\sum_{m,n=0}^{\infty}\frac{s^m t^n}{m!n!}H_{m,n}\left(x,y\right)H_{m,n}\left(\alpha,\beta\right) = \frac{1}{1-st}\exp\left[\frac{sx\alpha+ty\beta-\left(xy+\alpha\beta\right)st}{1-st}\right], \tag{E3}$$

Eq.(E2) can be rewritten as

$$(\text{E2}) = \frac{e^{-2|\alpha|^2}}{2T}e^{\frac{\frac{4e^{-\gamma t}}{\sqrt{T+1}}\left(\alpha^*\beta+\alpha\beta^*\right)-\left(\alpha^*\alpha+\beta\beta^*\right)\frac{4e^{-2\gamma t}}{T+1}}{1-\frac{e^{-2\gamma t}}{T+1}}}. \tag{E4}$$

Thus Eq.(25) becomes

$$
\begin{aligned}
W\left(\alpha,\alpha^{*},t\right) &= 4\int\frac{\mathtt{d}^{2}\beta}{\pi}e^{2\frac{T-1}{T+1}|\beta|^{2}}\sum_{m,n=0}^{\infty}C_{m,n}\left(\alpha,\alpha^{*},t\right)H_{m,n}\left(\frac{2\beta}{\sqrt{T+1}},\frac{2\beta^{*}}{\sqrt{T+1}}\right)W\left(\beta,\beta^{*},0\right)\\
&= \frac{2}{T}e^{-2|\alpha|^{2}}\int\frac{\mathtt{d}^{2}\beta}{\pi}\exp\left[2\frac{T-1}{T+1}\left|\beta\right|^{2}+\frac{2e^{-\gamma t}\left(\alpha^{*}\beta+\alpha\beta^{*}\right)-2e^{-2\gamma t}\left(\left|\alpha\right|^{2}+\frac{|\beta|^{2}}{T+1}\right)}{T}\right]W\left(\beta,\beta^{*},0\right)\\
&= R.H.S.\text{ of Eq.(28).}
\end{aligned}
\tag{E5}
$$

Appendix F: Derivation of Eq.(33)

Using the relation between Hermite polynomial and Lagurre polynomial,

$$
L_{m}\left(xy\right)=\frac{\left(-1\right)^{m}}{m!}H_{m,m}\left(x,y\right), \tag{F1}
$$

we can recast the left of Eq.(**33**) into the following form

$$
\begin{aligned}
F_{m,n} &= \frac{1}{s!}\int\frac{\mathtt{d}^{2}\alpha}{\pi}e^{-4|\alpha|^{2}}(-1)^{s}s!L_{s}\left(4\left|\alpha\right|^{2}\right)H_{m,n}\left(2\alpha^{*},2\alpha\right)\\
&= \frac{1}{s!}\int\frac{\mathtt{d}^{2}\alpha}{\pi}e^{-4|\alpha|^{2}}H_{s,s}\left(2\alpha^{*},2\alpha\right)H_{m,n}\left(2\alpha^{*},2\alpha\right)\\
&= \frac{1/4}{s!}\int\frac{\mathtt{d}^{2}\alpha}{\pi}e^{-|\alpha|^{2}}H_{s,s}\left(\alpha^{*},\alpha\right)H_{m,n}\left(\alpha^{*},\alpha\right).
\end{aligned}
\tag{F2}
$$

Further using the generating function of $H_{m,n}\left(\epsilon,\varepsilon\right)$,

$$
H_{m,n}\left(\epsilon,\varepsilon\right)=\frac{\partial^{m+n}}{\partial t^{m}\partial t'^{n}}\exp\left[-tt'+\epsilon t+\varepsilon t'\right]\Big|_{t=t'=0}, \tag{F3}
$$

and the integration formula,

$$
\int\frac{d^{2}z}{\pi}e^{\zeta|z|^{2}+\xi z+\eta z^{*}}=-\frac{1}{\zeta}e^{-\frac{\xi\eta}{\zeta}},\mathrm{Re}\left(\zeta\right)<0, \tag{F4}
$$

we have

$$
\begin{aligned}
&\int\frac{\mathtt{d}^{2}\alpha}{\pi}e^{-|\alpha|^{2}}H_{m',n'}\left(\alpha^{*},\alpha\right)H_{m,n}\left(\alpha^{*},\alpha\right)\\
&= \frac{\partial^{m+n}}{\partial t^{m}\partial t'^{n}}\frac{\partial^{m+n}}{\partial\tau^{m}\partial\tau'^{n}}\exp\left[-tt'-\tau\tau'\right]\\
&\times\int\frac{\mathtt{d}^{2}\alpha}{\pi}\exp\left[-\left|\alpha\right|^{2}+\left(t+\tau\right)\alpha^{*}+\left(t'+\tau'\right)\alpha\right]\Big|_{t=t'=\tau=\tau'=0}\\
&= \frac{\partial^{m'+n'}}{\partial t^{m'}\partial t'^{n'}}\frac{\partial^{m+n}}{\partial\tau^{m}\partial\tau'^{n}}\exp\left[t\tau'+\tau t'\right]_{t=t'=\tau=\tau'=0}\\
&= m!n!\delta_{m',n}\delta_{n',m},
\end{aligned}
\tag{F5}
$$

thus Eq.(F2) becomes the right hand side of Eq.(**33**).

Appendix G: Derivation of Eq.(35)

For this purpose, from Eq.(27) we have

$$
E_{m,n}=4\int\frac{\mathtt{d}^{2}\beta}{\pi^{2}}e^{-2y|\beta|^{2}}H_{m,n}\left(2x\beta,2x\beta^{*}\right)e^{-2|\beta-z|^{2}}, \tag{G1}
$$

where we have set

$$
y=\frac{1-\Lambda_{m,n}}{1+\Lambda_{m,n}},x=\frac{1}{\sqrt{\Lambda_{m,n}+1}}. \tag{G2}
$$

Using Eqs.(F3) and (F4), Eq.(G1) can be recast into the following form,

$$\begin{aligned} E_{m,n} &= 4e^{-2|z|^2}\frac{\partial^{m+n}}{\partial t^m\partial t'^n}e^{-tt'}\int\frac{\mathtt{d}^2\beta}{\pi^2}\exp\left[-2\left(y+1\right)\left|\beta\right|^2+2\beta\left(xt+z^*\right)+2\beta^*\left(xt'+z\right)\right]\Big|_{t=t'=0} \\ &= e^{-2|z|^2}\frac{\partial^{m+n}}{\partial t^m\partial t'^n}e^{-tt'}\int\frac{\mathtt{d}^2\beta}{\pi^2}\exp\left[-\frac{y+1}{2}\left|\beta\right|^2+\beta\left(xt+z^*\right)+\beta^*\left(xt'+z\right)\right]\Big|_{t=t'=0} \\ &= \frac{1}{\pi}\frac{2}{y+1}e^{-\frac{2y}{y+1}|z|^2}\frac{\partial^{m+n}}{\partial t^m\partial t'^n}\exp\left[-\left(1-\frac{2x^2}{y+1}\right)tt'+\frac{2xz}{y+1}t+\frac{2xz^*}{y+1}t'\right]_{t=t'=0}. \end{aligned} \tag{G3}$$

Note that $1-\frac{2x^2}{y+1}=0, y+1=\frac{1-\Lambda_{m,n}}{1+\Lambda_{m,n}}+1=\frac{2}{\Lambda_{m,n}+1}, -\frac{2y}{y+1}=\Lambda_{m,n}-1$, then we find

$$\begin{aligned} E_{m,n} &= \frac{1}{\pi}\frac{2}{y+1}e^{-\frac{2y}{y+1}|z|^2}\left(\frac{2xz}{y+1}\right)^m\left(\frac{2xz^*}{y+1}\right)^n \\ &= \frac{1}{\pi}\left(\Lambda_{m,n}+1\right)^{\frac{m+n+2}{2}}e^{(\Lambda_{m,n}-1)|z|^2}z^m z^{*n}. \end{aligned} \tag{G4}$$

[1] D. Bouwmeester et al., The Physics of Quantum Information, Springer, Berlin, 2000.

[2] E. Wigner, *Phys. Rev.* **40,** 749 (1932).

[3] Wolfgang P. Schleich, *Quantum Optics in Phase Space*, (Wiley-VCH, Birlin, 2001)

[4] A. Kenfack and K. Zyczkowski, J. Opt. B: Quantum Semiclass. Opt. **6**, 396 (2004).

[5] M. S. Kim and V. Bužek, Phys. Rev. A **46**, 4239 (1992).

[6] Li-yun Hu and Hong-yi Fan, J. Opt. Soc. Am. B, **25**, 1955 (2008); Li-yun Hu, Xue-xiang Xu and Hong-yi Fan, J. Opt. Soc. Am. B, **27**, 286 (2010).

[7] A. Biswas and G. S. Agarwal, Phys. Rev. A **75**, 032104 (2007).

[8] H. Jeong, A. P. Lund, and T. C. Ralph, Phys. Rev. A **72**, 013801 (2005).

[9] J. S. Neergaard-Nielsen, B. Melholt Nielsen, C. Hettich, K. Mølmer, and E. S. Polzik, Phys. Rev. Lett. **97**, 083604 (2006).

[10] A. Ourjoumtsev, A. Dantan, R. Tualle-Brouri, and Ph. Grangier, Phys. Rev. Lett. **98**, 030502 (2007).

[11] A. Ourjoumtsev, R. Tualle-Brouri, and Ph. Grangier, Phys. Rev. Lett. **96**, 213601 (2006).

[12] H. Jeong, J. Lee and H. Nha, J. Opt. Soc. Am. B, **25**, 1025 (2008).

[13] M. Stobinska, G. J. Milburn, and K. Wodkiewicz, Phys. Rev. A **78**, 013810 (2008).

[14] Hong-yi Fan and Li-yun Hu, Mod. Phys. Lett. B, **22**, 2435 (2008).

[15] Li-yun Hu and Hong-yi Fan, Opt. Commun. **282**, 4379 (2009); Hong-yi Fan and Li-yun Hu, Opt. Commun. **282,** 932 (2009); **281,** 5571 (2008).

[16] Memorial Issue for H. Umezawa, Int. J. Mod. Phys. B **10**, 1695 (1996) memorial issue and references therein.

[17] H. Umezawa, *Advanced Field Theory – Micro, Macro, and Thermal Physics* (AIP 1993).

[18] Y. Takahashi and H. Umezawa, Collecive Phenomena **2**, 55 (1975).

[19] Hong-yi Fan and J. R. Klauder, Phys. Rev. A **49,** 704 (1994).

[20] Li-yun Hu, Xue-xiang Xu and Shan-jun Ma, Int. J. Theor. Phys. (2010) DOI 10.1007/s10773-010-0434-y

[21] Hong-yi Fan and Yue Fan, J. Phys. A **35,** 6873 (2002); Hong-yi Fan and Hai-liang Lu, Mod. Phys. Lett. B, **21,** 183 (2007).

[22] Hong-yi Fan, Hai-liang Lu and Yue Fan, Ann. Phys. **321,** 480 (2006).

[23] Hong-yi Fan, H. R. Zaidi and J. R. Klauder, Phys. Rev. D **35**, 1831 (1987).

[24] A. Wünsche, J. Opt. B: Quantum Semiclass. Opt. **1,** R11 (1999).

[25] G. J. Milburn and C. A. Holmes, Phys. Rev. Lett. **56,** 2237 (1986); D. J. Daniel and G. J. Milburn, Phys. Rev. A **39** 4628 (1989).

[26] S. Chaturvedi, V. Srinivasan and G. S. Agarwal, J. Phys. A. Math. Gen **32,** (1999) 1909.

[27] G. J. Milburn and C. A. Holmes, Phys. Rev. A **44**, 4704 (1991)

[28] S. Chaturvedi and V. Srinivasan Phys. Rev. A **43**, 4054 (1991); S. Chaturvedi and V. Srinivasan J. Mod. Opt. **38**, 777 (1991).

[29] Li-yun Hu, Hong-yi Fan, Int. J. Theor. Phys. **48**, 3396 (2009).

[30] Hong-yi Fan, *Representation and Transformation Theory in Quantum Mechanics*, (Shanghai Scientific & Technical, Shanghai, 1997) (in Chinese).

[31] Hong-yi Fan and H. R. Zaidi, Phys. Lett. A **124**, 303 (1987).

[32] A. Wünsche, J. Computational and Appl. Math. **133**, 665 (2001).

[33] A. Wünsche, J. Phys. A: Math. and Gen. **33**, 1603 (2000).

[34] W. H. Louisell, *Quantum Statistical Properties of Radiation* (Wiley, New York, 1973).

[35] Hong-yi Fan and Li-yun Hu, Opt. Lett. **33,** (2008) 443.

[36] V. Perinova and A. Luks, Phys. Rev. A **41**, 414 (1990).